

SCOAP Rectal and Diverticulitis Surgery Dictation Guide

For rectal cancer:

- 1) Was procedure done for palliation? (Y/N)
- 2) Was preop (neoadjuvant) tx used? (Y/N)
 - a. If yes, was RT used? (Y/N)
 - b. If yes, was chemotherapy used? (Y/N)
- 3) Was preop fecal continence adequate? (Y/N)
- 4) Weeks between end of preop RT and surgery: (specify weeks)
- 5) Was EUS TRUS or MRI used to define stage? (Y/N)
- 6) What was the T stage (if known)? N stage? M stage?
- 7) Was distance of tumor from the anal verge defined? (Y/N)
 - a. If yes, was the distance determined by rigid scope, flexible scope and/or digital exam? (specify which)
 - b. If yes, was the distance determined after neoadjuvant therapy? (Y/N)
 - c. If yes, distance of tumor from anal verge: (specify cm)
- 8) Was tumor fixed to underlying structures: (Y/N)
 - a. If yes, was it fixed after neoadjuvant therapy? (Y/N)
- 9) Was a total mesorectal excision performed? (Y/N)
- 10) Was TME capsule intact? (Y/N)
- 11) Was a protective stoma used? (Y/N)
 - a. If no, was the anastomosis tested for leak with air or dye installation? (Y/N, specify method of testing)

For diverticular disease:

- 1) If colectomy for diverticular disease, was it for
 - a. ongoing, acute diverticulitis?
 - b. current GI bleeding?
 - c. colovesical fistula?
 - d. colonic stricture?
- 2) Did patient have prior episodes of diverticular disease (Y/N)
 - a. If yes, how many prior episodes of treated diverticulitis?
 - b. If yes, was patient treated as an inpatient for any of the episodes? (Y/N)